

A Note on the Weak Isovariant Borsuk-Ulam Theorem

Ikumitsu NAGASAKI ¹⁾

Abstract. In this note, we provide a refinement on the weak isovariant Borsuk-Ulam theorem. The main result is the following: For an arbitrary compact Lie group G , there exists a positive constant $c_G > 0$ such that, for any G -representations V and W , if there exists a G -isovariant map $f: V \rightarrow W$, then the inequality $c_G(\dim V - \dim V^G) \leq \dim W - \dim W^G$ holds.

1. The weak isovariant Borsuk-Ulam theorem

Let G be a compact Lie group. A G -equivariant map $f: X \rightarrow Y$ is called G -isovariant if f preserves the isotropy subgroups: $G_{f(x)} = G_x$, $x \in X$. In this note, all maps between spaces are continuous.

In [3], we have shown the following Borsuk-Ulam type result.

Theorem 1.1. *For an arbitrary compact Lie group G , there exists a weakly monotonely increasing function*

$$\varphi_G: \mathbb{N}_0 \rightarrow \mathbb{N}_0 \quad (\mathbb{N}_0 = \mathbb{N} \cup \{0\})$$

diverging to ∞ with the following property:

(P) *For any G -representations V and W , if there is a G -isovariant map $f: V \rightarrow W$, then*

$$\varphi_G(\dim V - \dim V^G) \leq \dim W - \dim W^G$$

holds.

However, a concrete form of φ_G is not given in [3]. In this note, we show that φ_G can be taken as a linear function $\varphi_G(n) = cn$ for some positive constant c . Namely, we show the following.

Dedicated to Professor Masakazu Tanatsugu on the occasion of his retirement from Kyoto Prefectural University of Medicine.

¹⁾ Department of Mathematics, Kyoto Prefectural University of Medicine, Shimogamohangi-cho 1-5, Sakyo-ku, Kyoto 606-0823, Japan. e-mail: nagasaki@koto.kpu-m.ac.jp

Theorem 1.2. *For an arbitrary compact Lie group G , there exists a positive constant $c > 0$ such that, for any G -representations V and W , if there exists a G -isovariant map $f: V \rightarrow W$, then*

$$c(\dim V - \dim V^G) \leq \dim W - \dim W^G$$

holds.

For a nontrivial compact Lie group G , we define \mathcal{C}_G as the set of $c \in \mathbb{R}$ such that

$$(1.1) \quad c(\dim V - \dim V^G) \leq \dim W - \dim W^G$$

holds for all G -representations V and W with a G -isovariant map $f: V \rightarrow W$.

Definition. We set $c_G = \sup \mathcal{C}_G$. For convenience, we set $c_G = 1$ when G is trivial.

Lemma 1.3. *The following hold.*

- (1) $0 \in \mathcal{C}_G$.
- (2) If $c' \leq c$ and $c \in \mathcal{C}_G$, then $c' \in \mathcal{C}_G$.
- (3) $0 \leq c_G \leq 1$.

Proof. (1) When $c = 0$, the inequality (1.1) clearly holds.

(2) This follows by $c'(\dim V - \dim V^G) \leq c(\dim V - \dim V^G) \leq \dim W - \dim W^G$.

(3) By (1), $0 \leq c_G$. Taking the identity map $id: V \rightarrow V$ for a nontrivial representation, we see that $c \leq 1$ by (1.1). Hence $c_G \leq 1$. \square

Lemma 1.4. *The supremum c_G is in the set \mathcal{C}_G ; i.e., c_G is the maximum of c satisfying inequality (1.1).*

Proof. If $c_G = 0$, then $c_G \in \mathcal{C}_G$ by Lemma 1.3 (1). Suppose that $c_G > 0$. For any $\varepsilon > 0$, $c_G - \varepsilon \in \mathcal{C}_G$ by Lemma 1.3 (2). Then we have

$$(c_G - \varepsilon)(\dim V - \dim V^G) \leq \dim W - \dim W^G \quad \text{for any } \varepsilon > 0.$$

This implies that $c_G(\dim V - \dim V^G) \leq \dim W - \dim W^G$. \square

Clearly, if $c_G = 1$ if and only if the isovariant Borsuk-Ulam theorem holds for G -representations; namely,

$$\dim V - \dim V^G \leq \dim W - \dim W^G$$

holds for all G -representations V and W with a G -isovariant map $f: V \rightarrow W$. The isovariant Borsuk-Ulam theorem was studied by Wasserman [6] and Nagasaki-Ushitaki

[5]. As a sufficient condition for $c_G = 1$, Wasserman gave the prime condition and Nagasaki and Ushitaki gave the Möbius condition. (For the details, see [5] and [6].) For example, from these conditions, one can see the following.

Proposition 1.5. *For the following groups, $c_G = 1$.*

- (1) *Solvable compact Lie group.*
- (2) *Alternating group A_n for $5 \leq n \leq 11$.*
- (3) *$\text{PSL}(2, q)$, q is a prime power.*

Remark. Some researchers conjecture that $c_G = 1$ for any finite group; however, it is still an open question.

2. Property of weak Borsuk-Ulam groups

We say that G is a *weak Borsuk-Ulam group* (weak BUG) if $c_G > 0$, and that G is a *Borsuk-Ulam group* (BUG) if $c_G = 1$. Theorem 1.2 implies that every compact Lie group G is a weak Borsuk-Ulam group, and Proposition 1.5 provides many examples of Borsuk-Ulam groups. We prepare several lemmas for showing Theorem 1.2.

Lemma 2.1. *Let H is a closed normal subgroup of G . If H and G/H is [weak] Borsuk-Ulam groups, then G is also a [weak] Borsuk-Ulam group.*

Proof. In the case where G and G/H are Borsuk-Ulam groups, Wasserman [6] has already shown it. Assume that G and G/H are weak Borsuk-Ulam groups. Let $f: V \rightarrow W$ be any G -isovariant map between representations. By restricting the action, $\text{res}_H f: V \rightarrow W$ is H -isovariant and by fixing by H , $f^H: V^H \rightarrow W^H$ is G/H -isovariant. By definition, we have

$$c_H(\dim V - \dim V^H) \leq \dim W - \dim W^H$$

and

$$c_{G/H}(\dim V^H - \dim V^G) \leq \dim W^H - \dim W^G.$$

By adding the inequalities, we obtain

$$c_H(\dim V - \dim V^H) + c_{G/H}(\dim V^H - \dim V^G) \leq \dim W - \dim W^G.$$

Setting

$$c = \min\{c_H, c_{G/H}\} > 0,$$

we obtain

$$c(\dim V - \dim V^G) \leq \dim W - \dim W^G.$$

This shows that $c_G > 0$. □

Lemma 2.2. *Let H be a closed normal subgroup of G . Then if G is a [weak] Borsuk-Ulam group, then G/H is also a [weak] Borsuk-Ulam group.*

Proof. In the case where G is a Borsuk-Ulam group, Wasserman [6] has already shown it. Assume that G is a weak Borsuk-Ulam group. Let $f: V \rightarrow W$ be a G/H -isovariant map between G/H -representations. By using the projection $p: G \rightarrow G/H$, V and W are thought of as G -representations and f is thought of as a G -isovariant map. Then, we have

$$c_G(\dim V - \dim V^G) \leq \dim W - \dim W^G.$$

Since $\dim V^G = \dim V^{G/H}$ and $\dim W^G = \dim W^{G/H}$, we have

$$c_G(\dim V - \dim V^{G/H}) \leq \dim W - \dim W^{G/H}.$$

This means that $0 < c_G \leq c_{G/H}$. □

Lemma 2.3. *Let H be a closed subgroup of G with $c_H = 1$. Assume that there exists a constant $0 < d < 1$ such that $\dim V^H \leq d \dim V$ for all nontrivial irreducible G -representations V . Then $c_G \geq 1 - d > 0$. In particular, G is a weak Borsuk-Ulam group.*

Proof. Let $f: V \rightarrow W$ be a G -isovariant map between G -representations. Decompose $V = V^\perp \oplus V^G$ and $W = W^\perp \oplus W^G$, where V^\perp [resp. W^\perp] denotes the orthogonal complement of V^G [resp. W^G]. Composing the natural inclusion $i: V^\perp \rightarrow V$ and the projection $p: W \rightarrow W^\perp$ with f , we obtain a G -isovariant map $g := p \circ f \circ i: V^\perp \rightarrow W^\perp$. Since H is a Borsuk-Ulam group, it follows that

$$\dim V^\perp - \dim V^{\perp H} \leq \dim W^\perp - \dim W^{\perp H} \leq \dim W^\perp.$$

By the complete reducibility of G , V^\perp decomposes into a direct sum of nontrivial irreducible representations. Hence by assumption one can see that

$$(1 - d) \dim V^\perp \leq \dim V^\perp - \dim V^{\perp H}.$$

Setting $c = 1 - d$, we obtain that $c \dim V^\perp \leq \dim W^\perp$, or equivalently

$$c(\dim V - \dim V^G) \leq \dim W - \dim W^G.$$

This implies that $c_G \geq c = 1 - d > 0$. □

Proposition 2.4. *A connected compact Lie group G is a weak Borsuk-Ulam group.*

Proof. Let T be a maximal torus of G . The following fact is proved in [3]. There exists a constant $0 < d < 1$ such that $\dim V^T \leq d \dim V$ for all nontrivial irreducible G -representations V . Therefore, by Lemma 2.3, a connected compact Lie group G is a weak Borsuk-Ulam group. \square

3. Proof of Theorem 1.2 and some examples

Let G be a (general) compact Lie group. Then there is a short exact sequence

$$1 \rightarrow G_0 \rightarrow G \rightarrow F \rightarrow 1,$$

where G_0 is the identity component of G and $F \cong G/G_0$ is a finite group. It follows from Proposition 2.4 that G_0 is a weak Borsuk-Ulam group. By Lemma 2.1, it suffices to show that a finite group is a weak Borsuk-Ulam group. By finite group theory, for every finite group G , there is a composition series

$$1 = H_0 \triangleleft H_1 \triangleleft \cdots \triangleleft H_r = G,$$

where each H_i/H_{i-1} is simple. Therefore, by using Lemma 2.1 repeatedly, it suffices to show that every finite simple group is a weak Borsuk-Ulam group. In [3], the following fact is proved.

Lemma 3.1. *Let G be a finite simple group. Let H be any nontrivial subgroup of G . Then there exists a constant $0 < d < 1$ such that $\dim V^H \leq d \dim V$ for all nontrivial irreducible representations V .*

In particular, if we take H as a nontrivial cyclic subgroup C , then since $c_C = 1$, it follows from Lemma 2.3 that every finite simple group is a weak Borsuk-Ulam group. As a consequence, an arbitrary finite group G is a weak Borsuk-Ulam group. Thus we obtain that every compact Lie group is a weak Borsuk-Ulam group.

Finally we give an example.

Proposition 3.2. *Let G be $\mathrm{SO}(3)$ or $\mathrm{SU}(2)$. Then $c_G \geq 4/5$.*

Proof. The fact $c_{\mathrm{SO}(3)} \geq 4/5$ has already shown in [4]. Indeed, if we take $\mathrm{O}(2) \subset \mathrm{SO}(3)$, then $\dim V^{\mathrm{O}(2)} \leq \frac{1}{5} \dim V$ for every nontrivial irreducible representation of $\mathrm{SO}(3)$. Since $\mathrm{O}(2)$ is solvable, $c_{\mathrm{O}(2)} = 1$. By Lemma 2.3, we have $c_{\mathrm{SO}(3)} \geq 4/5$.

Next we consider the case of $\mathrm{SU}(2)$. Let

$$T = \left\{ g_t := \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \mid t \in \mathbb{C}, |t| = 1 \right\}$$

be a maximal torus of $\mathrm{SU}(2)$. Let N be the normalizer of T in $\mathrm{SU}(2)$, which is isomorphic to $\mathrm{Pin}(2)$. Indeed, one can see

$$N = T \amalg Tb, \quad b = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

By representation theory (see [2] for example), $\mathrm{SU}(2)$ has one k -dimensional complex irreducible representation V_k for each $k \geq 1$. Let χ_k be the character of V_k . By the Weyl character formula,

$$\chi_k(g_t) = \begin{cases} t^{k-1} + t^{k-3} + \cdots + t^2 + 1 + t^{-2} + t^{-5} + \cdots + t^{-(k-1)} & \text{if } k \text{ is odd} \\ t^{k-1} + t^{k-3} + \cdots + t + t^{-1} + t^{-3} + \cdots + t^{-(k-1)} & \text{if } k \text{ is even} \end{cases}$$

for $g_t \in T$. Note that

$$\begin{aligned} \dim_{\mathbb{C}} V_k^T &= \int_T \chi_k(g_t) dt \\ &= \begin{cases} 1 & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even.} \end{cases} \end{aligned}$$

On the other hand, the order of g_tb is 4 and $(g_tb)^2 = -I = g_{-1} \in T$. Since g_tb is conjugate to g_i , we have

$$\chi_k(g_tb) = \chi_k(g_i) = \begin{cases} 1 & \text{if } k \equiv 1 \pmod{4} \\ -1 & \text{if } k \equiv -1 \pmod{4} \\ 0 & \text{otherwise} \end{cases}$$

and so

$$\begin{aligned} \dim_{\mathbb{C}} V_k^N &= \frac{1}{2} \left(\int_T \chi_k(g_t) dt + \int_{Tb} \chi_k(g_tb) dt \right) \\ &= \begin{cases} 1 & \text{if } k \equiv 1 \pmod{4} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus we have $\dim_{\mathbb{C}} V_k^N \leq \frac{1}{5} \dim_{\mathbb{C}} V_k$ for any k and so $\dim_{\mathbb{C}} V^N \leq \frac{1}{5} \dim_{\mathbb{C}} V$ for any complex representation V with $V^{\mathrm{SU}(2)} = 0$. For a real representation U , $U \oplus U$ has a complex structure. Then clearly $\dim_{\mathbb{C}} U \oplus U = \dim U$. Therefore we see that $\dim U^N \leq \frac{1}{5} \dim U$ for any real representation U with $U^{\mathrm{SU}(2)} = 0$. Since N is solvable, $c_N = 1$. By Lemma 2.3, we have $c_{\mathrm{SU}(2)} \geq 4/5$. \square

Remark. Determining the precise value of $c_{\mathrm{SU}(2)}$ or $c_{\mathrm{SO}(3)}$ is still an open question.

References

- [1] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, 1972.
- [2] T. Bröcker and T. tom Dieck, *Representations of compact Lie groups*, Springer, 1985.
- [3] I. Nagasaki, *The weak isovariant Borsuk-Ulam theorem for compact Lie groups*, Arch. Math. **81** (2003), 348–359.
- [4] I. Nagasaki, *Remarks on isovariant maps for representations*, RIMS Kôkyûroku **1343** (2003), 54–60.
- [5] I. Nagasaki and F. Ushitaki, *New examples of the Borsuk-Ulam groups*, RIMS Kôkyûroku Bessatsu **B39** (2013), 109–119.
- [6] A. G. Wasserman, *Isovariant maps and the Borsuk-Ulam theorem*, Topology Appl. **38** (1991), 155–161.